



South Asian Journal of Agricultural Sciences

E-ISSN: 2788-9297
P-ISSN: 2788-9289
SAJAS 2021; 1(2): 72-73
Received: 05-05-2021
Accepted: 07-06-2021

Karnewar SD
Assistant Professor,
College of Agril, Vidya
Pratishthan, Baramati,
Maharashtra, India

Dhawale RN
Assistant Professor,
VDCOAB, VNMKV,
Parbhani, Maharashtra, India

Dudhare MS
Associate Professor,
V.D. College of Agriculture
Biotechnology, VNMKV,
Parbhani, Maharashtra, India

Batish VK
Professor, National Dairy
Research Institute, Haryana,
India

Correspondence
Karnewar SD
Assistant Professor,
College of Agril, Vidya
Pratishthan, Baramati,
Maharashtra, India

Identification of lactobacilli at genus level by using PCR

Karnewar SD, Dhawale RN, Dudhare MS and Batish VK

Abstract

Lactic acid bacteria (LAB) are gram positive non sporulating, Microaerophilic bacteria whose main fermentation product from Carbohydrate is lactates. The LAB especially the genera Sactococcus, Lactobacillus, Leuconostoc, Streptococcus, Bifidobacteria and Pediococcus, Have been traditionally used as starter cultures for the fermentation of foods and beverages because of their contribution through flavor and aroma development and to spoilage retardation. Lactobacilli are found in milk, although their numbers are few and in certain cases even their absence has also been reported (Hill and Thornton, 1958) [3]. It is very difficult to isolate lactobacilli from milk as they are outnumbered by other microorganisms. Thus rendering most of the normal nutrient media unsuitable for their isolation. With the advent of molecular biology techniques, it is now possible to classify bacteria by comparing their genomes. Genotyping has many advantages over traditional typing procedures The rRNA locus is a genetic unit of broad evolutionary interest (Cedergren *et al.* 1988). In prokaryotes two additional assumptions are basic for the validity of this approach, namely that lateral gene transfer has not occurred between rRNA genes and that the amount of evolution or dissimilarity between rRNA sequences of a given pair of organisms is representative of the variation by the corresponding genomes (Rochelle, 1992) [7].

In present investigation 25 isolates alleged to be lactobacilli were recovered from BCP-MRS agar plates streaked with milk samples which was collected from NDRI, Karnal campus. These isolates on the basis of Grams staining and catalase test were subjected to PCR by using Lb1 and Lb2 23s rRNA lactobacilli specific PCR primers. Till the isolates produce a PCR amplified product of 194 bp specific for lactobacilli species on agarose gel which are consistent with the earlier observations of Suja (2003) [9].

Keywords: Lactobacilli, lactic acid bacteria, carbohydrate

Introduction

Lactobacilli are gram positive, catalase negative, non-spore forming rod shaped bacteria that produce lactic acid as the major end product of fermentation they have complex nutritional requirements regarding to be supplied with carbohydrates fatty acids or fatty acid esters. Salts nucleic acid derivatives and vitamins lactobacilli are found in milk although their numbers few and in certain cases even their absence has also been reported (Hill and thronton, 1958) [3]. It is very difficult to isolate lactobacilli from as they are outnumbered by other organisms, thus rendering most of the normal nutrient media unsuitable for their isolation (Chalmers, 1962) [2].

Lactobacilli constitute an important group of organisms which considerable improve the nutritional and therapeutic value of fermented foods. These organisms play an instrumental role in numerous fermentation processes in a variety of foods including dairy products, baked products and alcoholic beverages. The economic importance of lactobacilli clearly demands the development of a new technology to identify lactobacilli at DNA level. With the advent of molecular biology techniques, it is now possible to classify bacteria by comparing their genomes. Genotyping has many advantages over traditional typing procedures which includes 1. DNA can always be extracted from bacteria so that all strains are type able. 2. Analytical strategies for the Genotyping methods are similar and can be applied to DNA of Any source 3. In general, genomic DNA is a stable characteristic and its composition is independent of cultural conditions or methods PF preparation and 4. It allows for statistical data analysis and is amenable to automation.

In present investigation we isolated lactobacilli from milk samples and all the samples were subjected to grams staining and catalase test, the purity of the isolates as a lactobacilli were ascertained by using PCR techniques.

Materials and Methods

In present study, milk samples from fresh raw milk were collected from various location of NDRI, Karnal campus the samples were collected in sterile 10 ml capacity screw capped glass tubes after taking appropriate precautions as recommended by ISI (ISI, 1977) [4].

One ml of raw milk were added into 9 ml of litmus milk which was incubated overnight at 37°C and plated on BCP-MRS agar. Two types of colonies were observed, yellow colonies were picked up from the plates after an incubation period 24 hrs at 37 °C. totals of 25 isolated were picked up into MRS broth and further purified by repeated sub culturing. The purity of all bacterial isolates was always ascertained prior to use by Grams staining, cell morphology and catalase test. These isolates as a lactobacilli were then identified by using Lb1 and Lb2 primers (23s rRNA) specific to lactobacilli. The genomic DNA from the test cultures grown in MRS broth was extracted by following the method of Pospiech and Neikmann (1995) [6] method.

The PCR assay basically involves these steps viz. setting up of PCR reaction initial denaturation 95°C for 5 min, denaturation at 94 °C for 30 sec, renaturation at 56°C for 1 min, synthesis at 72°C temp for 1 min and extension at 72°C for 10 min by using MgCl₂, dNTPs 0.2ul, primer Lb1 and Lb2 2mM, taq DNA polymerase 2.25 units and template DNA about 100 ng to 1µg. The PCR amplified products obtained with different templates were electrophoresed on 2% agarose gel and examined under UV transilluminator (302 nm, photo dyne) and photographed using a Polaroid camera with Polaroid type 55 sheet film.


Results and Discussion

The role of LAB in human nutrition and health is well documented as those food grade bacteria have traditionally been used for various purposes. Among these LAB lactobacilli enjoy a special status, on account of their extensive use in food pharmaceutical and dairy industry as well. The commercial value of lactobacilli can be enhanced considerably by their genetic manipulation. The prerequisite for applying gene cloning technique to lactobacilli, is the development of method to identify lactobacilli at genus level (Vaishali *et al*, 1992) [10].

Lactobacilli are found in milk, although their numbers are few and in certain cases even their absence has also been reported (Hill and Thornton, 1958) [3] In present study, 25 isolates alleged to be lactobacilli were recovered from MRS agar plates streaked with milk samples pre enriched for 24 hrs in litmus milk and MRS broth for 24 hrs. Milton and Sandin (1986) [5] also developed a selective medium for Gram positive bacteria which was prepared using Elliker's lactic agar as the basal medium supplemented with 1% Tween 80.

The identity of the isolates which was recovered from yellow colonies as a lactobacilli was determined by Gram staining and then ascertained by PCR by using Lb1 and Lb2 set of primers specific to lactobacilli. All the isolates produce a PCR amplified product of 194 bp specific for lactobacillus species on agarose gel. The rRNA genetic locus is a unit of broad evolutionary interest in prokaryotes (Cedergren *et al*, 1988) [1]. Two additional assumptions are basic for the validity of this approach, namely that lateral gene transfer has occurred between rRNA genes and that the amount of evolution or dissimilarity between rRNA sequences of a given pair of organism is a representative of

the variations by corresponding genomes (Rochelle, 1992) [7]. Constituent results of 194 bp PCR product (specific for lactobacillus species on agarose gel), indicates that an important feature of the 16 sRNA molecule in its use as a universal standard parameter for phylogenetic interferences is the relative ease of sequence alignment. Thus, a myriad studies have exploited the features of 16 sRNA for speciation identification of lactobacilli at molecular level. It is evident that the resolving power of 16 sRNA sequences is limited when closely related species are being inspected.



References

1. Cedergren RJ, Gray MW, Abel Y, Sakoff D. The evolutionary relationship among known life forms. *J. of Molecular Evolution* 1988;28:98-112.
2. Chalmers CH. *Bacteria in relation to milk supply*, fourth edn, Edward Arnold Ltd. London 1962, 1-12.
3. Hill DA, Thornton HR. Lactobacilli in Edmonton dairy products. *Can J. Microbiol* 1958;4(3):215-220.
4. ISI, *Methods of test for dairy industry (Part 1), Rapid examination of milk (Part 3)* 1977, ISI: 1479.
5. Malton ME, Sandine WE. Improved media for differentiation of rods and cocci in Yoghurt. *J Dairy Sci* 1986;69:2569-2576.
6. Pospiech A, Neikmann B. A versatile quick preparation of genomic DNA from gram positive Bacteria. *TIG* 1997;11:217-218.
7. Rochelle AA, Fry JC, Weighmann AJ. DNA extraction for 16sRNA gene analysis to determine genetic diversity in deep sediment community. *FEMS microbiology Lett* 1992;100:59-65.
8. Rogosa DM, Sharpe EE. A medium for the cultivation of lactobacilli. *J. Appl. Bacteriol* 1960;23:130-235.
9. Suja. Development of PCR based techniques for identification of dairy lactobacilli, M. Sc thesis, NDRI, Karnal 2003.
10. Vaishali S, Batish VK, Grover S. Evidence of plasmid linked phenotypic characters in two wild lactobacilli casei strains, *Microbiologie-Aliments-Nutrition* 1992;10:169-175.