



## South Asian Journal of Agricultural Sciences

E-ISSN: xxxx-xxxx

P-ISSN: xxxx-xxxx

SAJAS 2021; 1(1): 38-42

Received: 12-11-2020

Accepted: 15-12-2020

**Ali H Jasim**

Faculty of Agriculture / Al-Qasim Al-Akhdar University, Babylon, Iraq

**Ali A Khalaf Arat**

Faculty of Agriculture / Al-Qasim Al-Akhdar University, Babylon, Iraq

# The effect of planting dates, plant density, and silicon spraying on the growth and yield of sweet corn

**Ali H Jasim and Ali A Khalaf Arat**

## Abstract

The experiment was carried out in private field in Abu Gharaq (10 km west of Hilla city) during the fall season 2019, to study the effect of planting dates (July 15, August 1 and 15) and plant density (40 and 80 thousand plants ha<sup>-1</sup>, silicon spraying levels (0, 1.5, 3.0, and 4.5 ml l<sup>-1</sup>) and their interactions on the growth and yield of sweet corn. Split-split-plot arrangement following a randomized complete block design (R.C.B.D) with three replications was used. The results were analyzed and the averages were tested according to the least significant difference at the 0.05 probability level. The most important results are summarized as follows: Planting on July 15 delayed the day's number to 50% silking to 53.21 days compared to August 1 (51.71 days), and decreased ripening single ear weight with sheaths to 104.3 g compared to August 15 which achieved 244.9 g. High plant density caused significant increase in 50% silking, and leaf area index to 53.75 days and 2.998, without significant effect on ripening single ear weight with sheaths. Silicon spraying levels caused significant effect on ear ripening weight, and 3 ml l<sup>-1</sup> achieved high weight 174.8 g compared to 169.4 g from control treatment.

**Keywords:** Sweet corn, Silicon, Planting date, Plant density

## Introduction

Sweet corn *Zea mays L. saccharata* is one type of yellow corn. Sweet corn is distinguished from yellow corn in containing its high sugar (14-20% sugar) seeds in the milky and early pasty stages (Rubatzky and Yamaguchi, 2000) [17]. Sweet corn has been cultivated in Iraq only as experimental scale, and due to the few studies, no field practices for its cultivation have been identified. Determining the appropriate date for planting is one of the important factors for crop growth and more efficient utilization of environmental factors that affect plant growth and yield (Azadbakht *et al.*, 2010) [7]. The plant density plays an important role in plant growth and yield, through its influence in determining the degree of competition between plants on light, water, and available nutrients (Van Roxel and Coulter, 2012) [21]. Silicon increased plant's ability to exploit solar energy and absorb light, which leads to increase the photosynthesis, and thus increase growth and yield (Deshmukh *et al.*, 2017) [9]. Silicon is one of the rare nutrients which takes an important role in bearing various types of stress and increases the rate of photosynthesis (Sahebi *et al.*, 2015) [18]. Silicon is present in the soil in large quantities, but it is one of the rare nutrients of the plant that it is preferred to add, since most silicon sources in the soil are in the form of crystals of insoluble aluminum silicates, which is not directly available to plants (Richmond and Sussman, 2003 and Meena *et al.*, 2014) [16, 14].

## Materials and methods

A field experiment was carried out in Abu Ghark district, within 32.3146 degrees north latitude and 44.2217 degrees east longitude, in the autumn season of 2019 to study the effect of planting dates (July 15, August 1 and 15), plant density (40 and 80 thousand plants per hectare) and silicon spraying levels (0, 1.5, 3, and 4.5 mL l<sup>-1</sup>) on growth and yield of sweet corn in silty clay loam soil (Table 1).

## Correspondence

**Ali H Jasim**

Faculty of Agriculture / Al-Qasim Al-Akhdar University, Babylon, Iraq

**Table 1:** Some physical and chemical field soil properties

characteristic	Value and unity	characteristic	Value and unity	
Available N	8.41 mg kg <sup>-1</sup>	Soil components	sand	186 g kg <sup>-1</sup>
Available P	19.02 mg kg <sup>-1</sup>		Silt	339 g kg <sup>-1</sup>
Available K	191 mg kg <sup>-1</sup>		clay	475 g kg <sup>-1</sup>
EC	2.9 dSm m <sup>-1</sup>	Soil texture		Silty clay loam
pH	7.5			

The experiment was carried as a factorial experiment according to split-split plots arrangement within a randomized complete block design with three replicates (3 \* 2 \* 4 \* 3). The main plots contained planting dated, and the split plots contained plant density, while silicon levels were taken place in split-split plots. Each experimental unit included three ridges, 1 m widths 3 m length, planted in one side, or both sides in hills 25 cm apart to attain the plant density. Compound fertilizer (N-P-K, 20-20-20) was added at 200 kg ha<sup>-1</sup> during soil preparation. The grain was seeded in the upper third of the ridge. The plants after two weeks thinned to one plant per hill. The data were analyzed and the means were tested according to the least significant differences at 0.05 probability.

**Results and discussion**

Table (2) shows that planting date had a significant effect on the number of days to 50% of tasselling, and planting on August 15 gave the highest number of days 43.12 days compared to July 15 that gave the lowest number 42.08 days. This may be due to day length high temperature during mid-July, which results in faster flowering (Oktem *et al.*, 2004) [15]. The plant density had a significant effect on this trait, and the low density gave the lowest day number 42.39 days compared to the high density (43.08 days). This

may be due to the deception that encouraged vegetative growth and delay of the transition period to reproductive growth. Or that increasing plant density led to a lack of light penetration inside the plant, that increasing ouxine and then prolonging the vegetative growth period and delaying the transition to reproductive status (Hashemi and Herbert, 1992) [10]. This result was agreed with Shrestha *et al* (2018) [20]. Silicon spraying had a significant effect and the level of 4.5, gave the highest days 43.33 days, while control treatment gave the lowest days number 42.22 days.

The interaction between the dates of planting and plant density had a significant effect and August 15 with high density gave the highest days number of days 43.50 days, while July 15 and the low density gave the lowest number 41.67 days. The interaction between planting date and silicon had a significant effect, and August 15 with silicon 4.5 ml l<sup>-1</sup> gave the highest days number 44.00 days, while July 15 without spraying silicon gave the lowest number (40.67 days). The interference between plant density and silicon had a significant effect, and 1.5 ml l<sup>-1</sup> silicon with high density gave the highest number 43.56 days compared to the low density without silicon (41.89 days). The interaction between the study factors had a significant effect and the lowest days number (40.33 days) achieved from July 15, low density, and without spraying silicon.

**Table 2:** Effect of seeding dates, plant density and Si spraying on tasselling (days)

Seeding dates	Plant Density	Silicon treatments				Interaction of dates* density
		0	1.5	3	4.5	
July 15	40000	40.33	42.33	41.33	42.67	41.67
	80000	41.00	44.00	42.67	42.33	42.50
August 1	40000	42.67	41.67	42.67	44.00	42.75
	80000	43.33	43.33	43.33	43.00	43.25
August 15	40000	42.67	41.67	42.67	44.00	42.75
	80000	43.33	43.33	43.33	44.00	43.50
Si mean		42.22	42.72	42.67	43.33	
LSD <sub>0.05</sub>		Si= 0.688 T=2.186				1.420
Interaction of dates * silicon						Dates average
July 15		40.67	43.17	42.00	42.50	42.08
August 1		43.00	42.50	43.00	43.50	43.00
August 15		43.00	42.50	43.00	44.00	43.12
LSD <sub>0.05</sub>		1.720				0.773
Interaction of density * silicon						Density average
40000		41.89	41.89	42.22	43.56	42.39
80000		42.56	43.56	43.11	43.11	43.08
LSD <sub>0.05</sub>		1.438				0.631

Table (3) shows that planting date had a significant effect on the number of days to 50% of silking, and planting on July 15 gave the highest number of days 53.21 days compared to August 1 that gave the lowest number 51.71 days. This may be due to high temperature during mid-July, which results in faster flowering (Baker and Ahmed, 2002) [8]. This result was agreed with Al-Zuhairy (2005) [6]. The plant density had a significant effect on this trait, and the high density gave the highest day number 53.75 days compared to the low density (51.44 days). This may be due to the deception

that increasing ouxine and then prolonging the vegetative growth period (Al-Taee, 2007) [5]. This result was agreed with Shrestha *et al* (2018) [20].

The interaction between the dates of planting and plant density had a significant effect and July 15 with high density gave the highest days number 55.25 days, while August 1 and the low density gave the lowest number 50.92 days. The interaction between planting date and silicon had a significant effect, and July 15 without silicon gave the highest days number 53.83 days, while August 1 without

spraying silicon gave the lowest number (51. days). The interaction between plant density and silicon had a significant effect, and 1.5 ml l<sup>-1</sup> silicon with high density gave the highest days 54.11 days compared to the low density without silicon (51.11 days). The interaction

between the study factors had a significant effect and the highest days number (56.00 days ) achieved from July 15, 1.5 ml l<sup>-1</sup> silicon, and high density, while August 1, low density, and without spraying silicon gave the lowest days 50.00 days.

**Table 3:** Effect of seeding dates, plant density and Si spraying on silking (days)

Seeding dates	Plant density	Silicon treatments				Interaction of dates* density
		0	1.5	3	4.5	
July 15	40000	52.00	51.33	51.00	50.33	51.17
	80000	55.67	56.00	54.33	55.00	55.25
August 1	40000	50.00	51.33	51.33	51.00	50.92
	80000	52.00	53.00	53.00	52.00	52.50
August 15	40000	51.33	52.00	52.67	53.00	52.25
	80000	52.33	53.33	54.00	54.33	53.50
Si mean		52.22	52.83	52.72	52.61	
LSD <sub>0.05</sub>		Si=n.s T=1.555				1.050
Interaction of dates * silicon						Dates average
July 15		53.83	53.67	52.67	52.67	53.21
August 1		51.00	52.17	52.17	51.50	51.71
August 15		51.83	52.67	53.33	53.67	52.88
LSD <sub>0.05</sub>		1.194				1.019
Interaction of density * silicon						Density average
40000		51.11	51.56	51.67	51.44	51.44
80000		53.33	54.11	53.78	53.78	53.75
LSD <sub>0.05</sub>		0.848				0.535

Table (4) shows that plant density caused a significant effect on the plant leaf area index, and high density gave the highest value 2.998, while low density gave the lowest value 1.494. This result was agreed with Abd Salama *et al.* (2007). The interaction between the dates of planting and plant density caused a significant effect, and August 1 with high density was superior by giving the highest value 3.068,

while the July 15 and low density gave the lowest value (1.442). The interaction between planting dates and silicon levels caused a significant effect, and August 1 without spraying silicon gave the highest value 2.372, compared to July 15 without spraying silicon, which gave the lowest value 2.095. The highest leaf area index (3.243) achieved from August 1 with high density and without silicon.

**Table 4:** Effect of seeding dates, plant density and Si spraying on leaf area index

Seeding dates	Plant density	Silicon treatments				Interaction of dates* density
		0	1.5	3	4.5	
July 15	40000	1.453	1.417	1.480	1.417	1.442
	80000	2.737	3.240	2.790	3.003	2.943
August 1	40000	1.500	1.463	1.613	1.417	1.498
	80000	3.243	3.083	2.870	3.073	3.068
August 15	40000	1.503	1.577	1.573	1.517	1.543
	80000	2.960	2.943	2.963	3.063	2.983
Si mean		2.233	2.287	2.215	2.248	
LSD <sub>0.05</sub>		Si=n.s T=0.241				0.162
Interaction of dates * silicon						Dates average
July 15		2.095	2.328	2.135	2.210	2.192

August 1	2.372	2.273	2.242	2.245	2.283
August 15	2.232	2.260	2.268	2.290	2.263
LSD <sub>0.05</sub>	0.187				n.s
Interaction of density * silicon					Density average
40000	1.486	1.486	1.556	1.450	1.494
80000	2.980	3.089	2.874	3.047	2.998
LSD <sub>0.05</sub>	0.129				0.076

Table (5) shows that planting dates caused a significant effect on the marketable ear weight with sheath (g), and August 15 was significantly distinguished by giving the highest weight 244.9 g, while July 15 gave the lowest weight 104.3 g. This result was due to the optimum temperatures during the flowering period that affected the fertilization process and increases ear grains number (data not publish), which is reflected positively in increasing yield (Ahmed, 2001 and Al-Hassani, 2015) <sup>[1, 3]</sup>. This is consistent with the findings of Al-Hadidi (2007) and Al-Mashhadani (2010) <sup>[2, 4]</sup>. Silicon spraying caused a significant effect and the level 3.0 ml l<sup>-1</sup> gave the highest weight of 183.3 g, while control treatment gave the lowest weight of 169.4 g. This is due to the role of silicon in improving the activity of photosynthesis and the efficiency of its representation in the plant, thereby increasing the dry matter (Jawahar and Vaiyapuri, 2010, 2013 and Xie *et al*, 2014) <sup>[12-13, 22]</sup>. These results are consistent with Zhiming *et al* (2014) and Idan (2016) <sup>[23, 11]</sup>.

The interaction between planting dates and plant density had a significant effect August with low density gave the highest weight 260.8, while July 15 with high density gave the

lowest weight 101.9. The interaction between planting dates and silicon levels caused a significant effect, and August 15 with 3 ml l<sup>-1</sup> silicon gave the highest weight 251.7 g, while July 15 at all-silicon spraying levels gave the lowest weight. The interaction between plant density and silicon spraying levels caused a significant effect and low density with 3.0 ml l<sup>-1</sup>silicon gave the highest weight 185.8 g, while the high density without silicon spray gave the lowest weight 164.8 g. The highest weight achieved from August 15 with low density and sprayed silicon at 1.5 - 4.5 ml l<sup>-1</sup> reached to 266.0 - 269.3 g, while July 15 with high density at all levels of silicon gave the lowest ripening ear weight with sheath.

**Conclusion**

It was concluded from this experiment, that delayed planting date of sweet corn to 15 August improved the weight of ripening single ear with sheaths. Planting high density 80 000 plants ha<sup>-1</sup> had no significant effect on the weight of ripening single ear with sheaths, but of course doubled the ear numbers per unit area. Silicon spraying caused an increase in the weight of ripening single ear with sheaths.

**Table No 5:** Effect of seeding dates, plant density and Si spraying on harvesting ear wt

Seeding dates	Plant density	Silicon treatments				Interaction of dates* density
		0	1.5	3	4.5	
July 15	40000	103.0	111.7	109.0	103.0	106.7
	80000	98.3	107.7	98.7	103.0	101.9
August 1	40000	179.3	183.7	182.3	176.7	180.5
	80000	175.3	171.3	206.0	164.0	179.3
August 15	40000	240.0	268.0	266.0	269.3	260.8
	80000	220.7	224.7	237.3	232.7	228.9
Si mean		169.4	181.2	183.3	174.8	
LSD <sub>0.05</sub>		Si= 9.42 T=24.8				17.2
Interaction of dates * silicon						Dates average
July 15		100.7	109.7	103.8	103.0	104.3
August 1		177.3	177.5	194.5	170.3	179.9
August 15		230.3	246.4	251.7	251.0	244.9
LSD <sub>0.05</sub>		19.8				17.4
Interaction of density * silicon						Density average
40000		174.1	181.1	185.8	176.3	179.3
80000		164.8	181.2	180.7	173.2	177.5
LSD <sub>0.05</sub>		12.8				n.s

**References**

1. Ahmed SA. The stages and characteristics of growth and yield of maize genotypes (*Zea mays* L.) by

effecting of planting dates. Master Thesis, College of Agriculture - University of Baghdad 2001.

2. Al-Hadidi KHK. Effect of sowing date and distance between lines on yield and its components for two maize varieties of maize. M.Sc. Thesis, College of Agriculture - University of Mosul 2007.
3. Al-Hassani MHO. Sweet corn response (*Zea mays* L. *saccharata*) for planting dates and spraying with bio-stimulants. PhD thesis - College of Agriculture - University of Baghdad 2015.
4. Al-Mashhadani NA. Effect of sowing dates on yield and its components for five genotypes of maize. *Anbar J Agric Sci* 2010;8(2):64-70.
5. Al-Taei ADJ. The effect of plant density and levels of nitrogen fertilizer on yield and some field traits of sweet corn. Master Thesis. Department of Field Crops. College of Agriculture, University of Baghdad 2007, 91-101.
6. Al-Zuhairi NSA. Estimation of genetic parameters in crossbreeds of maize (*Zea mays* L.). Master Thesis, College of Agriculture and Forestry - University of Mosul 2005.
7. Azadbakht AG, Azadbakht H, Nasrollahi and Z Bitarafan. Evaluation of Different planting Dates Effect on Three Maize Hybrids in Koohdasht Region of Iran. *Int. J. Sci. and Adv. Tech* 2010;2(3):34-38.
8. Bakr RH and Ahmed SA. The yield and its components of maize grown on different dates. *Iraqi Agricultural Science Journal* 2002;33(5):121-130.
9. Deshmukh RK, J Ma and RR Belanger. Role of silicon in plants. *Front Plant Sci* 2017;8:1858.
10. Hashemi A and SJ Herbert. Intensifying plant density response of corn with artificial shade. *Agron J* 1992;84:547-551.
11. Idan Sarmad Faisal. Effect of Etherel and Silicon Interaction on growth and yield of Popcorn (*Zea mays ssp.everta*) under Deficit Irrigation. M.Sc. Thesis, Agric. Coll., Al-Qasim Green Univ 2016.
12. Jawahar S and Vaiyapuri V. Effect of sulphur and silicon fertilization on growth and yield of rice. *Int. J. Current Res* 2010;9(1):36-30.
13. Jawahar S and Vaiyapuri V. Effect of sulphur and silicon fertilization on yield, nutrient uptake and economics of rice. *Int. Res. J. Chem* 2013;1(1):34-43.
14. Meena VD, ML Dotaniya, V Coumar, S Rajendiran, S Kundu, AS Rao. A case for silicon fertilization to improve crop yields in tropical soils. *Proc. Natl. Acad. Sci., India, Sect. B Biol. Sci* 2014;84(3):505-518.
15. Oktem A, AG Oktem And Y Coskun. Determination of sowing dates of sweet corn (*Zea mays* L. *saccharata* *Sturt.*) under Sanliurfa Conditions. *Turk. J. Agric. & For* 2004;28(2):83-91.
16. Richmond KE and Sussman M. Got silicon The non-essential beneficial plant nutrient. *Current Opinion in Plant Biol* 2003;6(3):268-272.
17. Rubatzky VE and M Yamaguchi. *World vegetables: principles, production, and nutritive values* (2<sup>nd</sup> ed.). New York: Chapman & Hall 2000.
18. Sahebi M Hanafi M, M Akmar, AS Rafii, MY Azizi, P Tengoua FF. Importance of silicon and mechanisms of bio-silica formation in plants. *Bio-Med Res, Int.* 1-16.
19. Salama MA, HB Aswad and HS Mahdi. Effect of space between plants and nitrogen fertilizer on growth and grain yield of corn. *Al-Anbar J. Agric. Sci* 2007;5(1):143-148.
20. Shrestha J, DN Yadav, LP Amgain and JP Sharma. Effects of nitrogen and plant density on maize (*Zea mays* L.) phenology and grain yield. *Current Agric. Res. J* 2018;6(2):175-182.
21. Van Roekel RJ and JA Coulter. Agronomic responses of corn hybrids to row width and plant density. *Agron. J* 2012;104:612-620.
22. Xie ZF, Song H Xu, H Shao and R Song. Effects of silicon on photosynthetic characteristics of maize (*Zea mays* L.) on alluvial soil. *The Sci. World J* 2014;14:1-6.
23. Zhiming X and Fengbin S. Effects of silicon on photosynthetic characteristics of maize (*Zea mays* L.) on alluvial soil. *Sci. World J* 2014;71(8):102-112.